

# ANASCORP®

Centruroides (Scorpion)  
Immune F(ab')<sub>2</sub> (Equine) Injection



## SYMPTOM RECOGNITION & DOSING REGIMEN

### Is Your Emergency Department Prepared?

Why send your patient to the ICU when you can send them home in under 4 hours?  
95-100% of patients experienced symptom relief in less than four hours after initiating ANASCORP treatment.\* Do your best to alleviate hardships on families due to hospital admissions, transfer costs and complications of supportive care by using ANASCORP.

**ANASCORP is the only FDA-approved antivenom indicated for treatment of clinical signs of scorpion envenomation.\*** Make sure your hospital knows how to recognize envenomation symptoms, when to use ANASCORP, and how to properly dose ANASCORP.

### SYMPTOM RECOGNITION

Symptoms can include: pain, loss of muscle control, roving or abnormal eye movements, slurred speech, respiratory distress, hypersalivation, airway obstruction and vomiting.<sup>1</sup> Symptoms are graded on a 4-point grading scale, adapted from O'Connor:<sup>1</sup>

#### MILD ENVENOMATION

**GRADE 1:** Localized pain and/or paresthesias at the site of envenomation.

**GRADE 2:** Pain and/or paresthesias, remote from site of envenomation.

#### SEVERE ENVENOMATION


Often requires emergency management in a healthcare setting

**GRADE 3:** **Either** cranial nerve dysfunction: Tongue fasciculations, hypersalivation, slurred speech or opsoclonus.  
**Or** neuromuscular dysfunction: Involuntary shaking and jerking of extremities.

**GRADE 4:** **Both** cranial nerve **AND** neuromuscular dysfunction.

Bark scorpion venom can affect people of all ages, but the majority of severe envenomations occur in children. Of the 1,534 patients in the clinical trials, 78% were children.\*

### ANASCORP DOSING REGIMEN\*



\*Please see accompanying package insert for full prescribing information.

†Sterile normal saline (0.9% NaCl).

<sup>1</sup>O'Connor AD, Ruha AM. Clinical course of bark scorpion envenomation managed without Antivenom. J Med Toxicol. 2012 Sep;8(3): 258-62.

# ANASCORP<sup>®</sup>

Centruroides (Scorpion)  
Immune F(ab')<sub>2</sub> (Equine) Injection

## IMPORTANT STOCKING CONSIDERATIONS

1. Initial dose is 3 vials.
2. Stocking recommendation is a minimum of 5 vials per patient. Extended holiday weekends may require having additional vials on hand.
3. ANASCORP is only available from AnovoRx: 1 (866) 830-743.7.
4. 3-year expiration dating.
5. Free replacement for expired, unopened vials. ‡



## SAFETY\*

### ADVERSE REACTIONS

The most common adverse reactions observed in ≥ 2% of patients in the clinical studies for ANASCORP were: vomiting, pyrexia, rash, nausea and pruritus.

### Postmarketing Experience

The following adverse reactions have been identified during post approval use of ANASCORP: chest tightness, palpitations, rash and pruritus.

### WARNINGS AND PRECAUTIONS

#### Hypersensitivity Reactions

Severe hypersensitivity reactions, including anaphylaxis, are possible with ANASCORP. Prepare for monitoring and management of allergic reactions, particularly in patients with a history of hypersensitivity to equine (horse) proteins or patients who have received previous therapy with antivenoms containing scorpion or equine proteins.

#### Delayed Allergic Reactions (Serum Sickness)

Delayed allergic reactions (serum sickness) may occur following treatment with ANASCORP. Patient monitoring with follow-up visit is recommended.

#### Transmissible Infectious Agents

ANASCORP is made from equine plasma and may contain infectious agents, e.g. viruses.

#### Reaction to Cresol

Localized reactions and generalized myalgias have been reported with the use of cresol as an injectable excipient.

## BE PREPARED

**Why send your patient to the ICU when you can send them home in under 4 hours?**

Make sure you are prepared for a bark scorpion sting emergency.

Do your best to alleviate hardships on families due to hospital admissions, transfer costs and complications of supportive care.

Use ANASCORP to treat clinical signs of scorpion envenomation in your Emergency Department.

To schedule an in-service or for training materials contact:

**Paul Landes**  
602-741-6077  
inservice@raretx.com

Training video at:  
[www.anascorp-us.com](http://www.anascorp-us.com)

**ANASCORP** is only available from:  
**AnovoRx: 866-830-7437**  
[www.anovorx.com](http://www.anovorx.com)

\*Please see accompanying package insert for full prescribing information.

† See Return Goods Policy for terms and conditions

© Rare Disease Therapeutics, Inc. | RDT Anascorp Cut-1-001 | September 2018